

TRADITIONAL HERBAL KNOWLEDGE AND MEDICINAL PLANT USE AMONG TRIBAL COMMUNITIES IN INDIA

Ms. Shilpa Gorakh Chavan¹, Dr. Bharat Bhushan²

Research Scholar, Department of Botany, Sunrise University, Alwar¹

Research Guide, Department of Botany, Sunrise University, Alwar²

ABSTRACT

Ethnobotany plays a pivotal role in documenting indigenous knowledge of medicinal plants among tribal communities in India, where biodiversity hotspots coincide with rich cultural traditions. This comprehensive review and meta-analysis synthesizes findings from over 150 ethnobotanical studies conducted across diverse tribal regions between 1995 and 2025. The primary objectives were to quantify trends in documentation efforts, identify commonly utilized plant parts, evaluate conservation strategies, and critically assess gaps in past research methodologies. Data were systematically extracted from peer-reviewed journals, including the Journal of Ethnopharmacology, Indian Journal of Traditional Knowledge, and Biodiversity and Conservation. Results indicate a steady increase in publications post-2010, with leaves (32%) and roots (22%) being the most frequently documented plant parts. Tribal communities in the Northeast, Central India, and Western Ghats regions exhibit the highest ethnobotanical activity. However, critical analysis reveals significant methodological inconsistencies, including inadequate sample sizes, lack of quantitative indices in earlier studies, and insufficient integration of molecular validation. Major threats identified through the meta-analysis include habitat loss (impact score 87), overharvesting (72), and policy gaps. Emerging conservation approaches such as community-based management and ex-situ cultivation show promise but remain under-adopted (overall adoption <60%). The findings underscore the urgent need for standardized protocols, digital databases, and participatory approaches that integrate traditional ecological knowledge with modern conservation science. This review provides a foundational framework for future ethnobotanical research and policy formulation aimed at sustainable utilization and conservation of India's medicinal plant heritage.

Keywords: *Ethnobotany¹, Medicinal plants², Tribal communities³, Conservation⁴, Documentation⁵, Meta-analysis⁶, Biodiversity hotspots⁷, Traditional knowledge⁸*

1. INTRODUCTION

1.1 Background and Significance of Ethnobotanical Research in India

India harbors one of the world's richest repositories of medicinal plants and indigenous knowledge systems, with over 8,000 species documented for therapeutic applications. Tribal communities, comprising approximately 8.6% of the national population and residing predominantly in forested and hilly terrains, have preserved sophisticated ethnobotanical practices spanning millennia. These communities, including the Bhils, Gonds, Santhals, Nagas, and Todas, rely heavily on plant-based remedies for primary healthcare due to limited access to modern medical infrastructure. Ethnobotanical documentation serves not only as a repository of traditional wisdom but also as a critical resource for pharmaceutical bioprospecting, biodiversity conservation, and cultural heritage preservation[1]. Past research has established that nearly 70% of the rural population in tribal belts continues to depend on herbal medicine. However, rapid socioeconomic transformations, deforestation, and climate change threaten both the plant resources and the transmission of this knowledge to younger generations. Meta-analyses of ethnobotanical literature are essential to synthesize fragmented studies, identify knowledge gaps, and guide evidence-based conservation policies. The integration of traditional ecological knowledge (TEK) with scientific validation offers immense potential for sustainable development goals and the protection of intellectual property rights under frameworks such as the Convention on Biological Diversity (CBD). This review addresses these imperatives by conducting a rigorous meta-analysis focused on past work pertaining to documentation and conservation efforts[2].

1.2 Objectives of the Present Meta-Analysis

The overarching aim of this review is to systematically evaluate and synthesize the body of past ethnobotanical research on medicinal plants utilized by tribal communities in India. Specific objectives include: (1) mapping the temporal and geographical distribution of documentation studies and quantifying publication trends from 1995 to 2025; (2) identifying predominant plant families, species, and plant parts employed in traditional medicine across major tribal regions; (3) assessing the adoption and effectiveness of various conservation strategies reported in the literature; (4) critically analyzing methodological strengths and limitations of prior studies to highlight best practices and recurring pitfalls; and (5) formulating evidence-based recommendations for future research and policy interventions[3]. By employing meta-analytic techniques, this study aggregates quantitative and qualitative data from diverse sources to derive robust conclusions that transcend individual case studies. The analysis also examines how factors such as accessibility, literacy levels, and government interventions influence the continuity of ethnobotanical knowledge. These objectives directly respond to calls from international bodies like the World Health Organization (WHO) and the International Union for Conservation of Nature (IUCN) for the integration of indigenous knowledge into national biodiversity strategies. Ultimately, the review seeks to bridge the gap between academic documentation and actionable conservation outcomes[4].

1.3 Scope, Geographical Coverage, and Structure of the Review

This meta-analysis encompasses ethnobotanical studies conducted across 28 Indian states with a particular emphasis on six major tribal belts: the Northeast (Assam, Nagaland, Manipur, Meghalaya), Central India (Chhattisgarh, Jharkhand, Odisha, Madhya Pradesh), Western Ghats (Kerala, Karnataka, Maharashtra), Himalayan foothills (Uttarakhand, Himachal Pradesh), Eastern Ghats, and the Aravalli ranges in Rajasthan and Gujarat. Studies published in English between 1995 and 2025 were included, focusing on peer-reviewed articles

that reported primary field data on medicinal plant use among recognized tribal groups. Excluded were purely pharmacological or phytochemical studies lacking ethnobotanical context, as well as reviews without original data synthesis[5]. The review is structured into eight main sections. Following the introduction, a comprehensive survey of past literature is presented, followed by a detailed methodology section outlining the meta-analytic protocol. A critical analysis evaluates methodological rigor and thematic gaps. Discussion integrates findings with broader ecological and socio-cultural contexts, leading to a concise conclusion and a set of actionable recommendations. Five original figures synthesized from aggregated data illustrate key patterns. The references comprise 30 sources in IEEE format drawn from leading ethnobotany and conservation journals. This structured approach ensures transparency, reproducibility, and utility for researchers, policymakers, and conservation practitioners alike[6].

2. SURVEY OF PAST WORK AND META-ANALYSIS OF DOCUMENTATION EFFORTS

The systematic survey of past ethnobotanical literature reveals a rich but heterogeneous body of research spanning three decades. Early foundational studies in the 1990s, such as those by Jain [1] and Pushpangadan [2], laid the groundwork by cataloging medicinal plants among the tribes of Central India and the Western Ghats. These pioneering works were primarily qualitative inventories that recorded vernacular names, therapeutic uses, and modes of preparation without employing quantitative indices. The period from 2000 to 2010 witnessed a surge in regional surveys, particularly in the biodiversity hotspots of Northeast India. Researchers documented over 300 species used by the Naga, Khasi, and Garo communities, with emphasis on plants for treating gastrointestinal, respiratory, and dermatological ailments [3], [4]. Meta-analytic aggregation indicates that during this decade, leaves constituted the dominant plant part (35-40%), followed by roots and bark. A landmark contribution came from the All India Coordinated Research Project on Ethnobiology (AICRPE), which compiled data across 24 states and identified more than 10,000 plant-based preparations [5]. However, these compilations suffered from inconsistent methodologies and limited statistical validation.

Post-2010 research demonstrated marked improvements in methodological sophistication. The introduction of quantitative ethnobotanical indices such as Use Value (UV), Informant Consensus Factor (ICF), and Fidelity Level (FL) became standard practice, enabling cross-study comparisons [6], [7]. Studies in Odisha and Chhattisgarh by researchers like Sahu et al. [8] and Sharma et al. [9] applied these indices to evaluate the cultural importance of species such as *Andrographis paniculata*, *Curcuma longa*, and *Rauvolfia serpentina* among tribal healers. The meta-analysis of 87 studies revealed that the average number of informants per study increased from 45 (pre-2010) to 112 (2015-2025), reflecting greater community engagement. Geographically, the Western Ghats and Northeast regions accounted for 48% of all documented species, while Central Indian studies contributed disproportionately to knowledge of anti-malarial and anti-diabetic plants. Publication trends analyzed in Figure 2 demonstrate a near-linear growth, with annual outputs rising from approximately 28 papers in the late 1990s to over 450 in recent years. This expansion correlates with increased funding from the National Medicinal Plants Board (NMPB) and the Department of Biotechnology (DBT).

A detailed examination of plant part utilization across aggregated data (Figure 3) shows leaves remain the most frequently cited (32%), followed by roots/rhizomes (22%), bark (15%), fruits/seeds (12%), flowers (8%), whole plants (7%), and stems (4%). This distribution aligns with global patterns and reflects ease of collection,

seasonal availability, and traditional preferences for non-destructive harvesting. Dominant families identified in the meta-analysis include Fabaceae (12.4%), Asteraceae (9.8%), Lamiaceae (8.1%), Euphorbiaceae (7.3%), and Apocynaceae (6.5%). Species with highest consensus across tribal groups include *Azadirachta indica*, *Ocimum sanctum*, *Tinospora cordifolia*, and *Withania somnifera*. Regional variations are pronounced: Himalayan tribes heavily utilize alpine herbs such as *Aconitum heterophyllum* and *Picrorhiza kurroo* for fever and liver disorders, whereas Western Ghats communities emphasize tree barks and climbers for rheumatism and snakebites [10], [11]. The Northeast stands out for its documentation of orchids and ferns in traditional healing. Despite this wealth of data, the survey highlights that fewer than 15% of documented plants have undergone rigorous pharmacological or clinical validation, representing a significant translational gap.

Conservation-oriented studies gained momentum after 2012, coinciding with India's ratification of the Nagoya Protocol. Research by Rao et al. [12] and Singh and Kumar [13] assessed the status of threatened medicinal plants in tribal areas using IUCN criteria and reported that 28% of frequently used species are vulnerable or endangered. Community-based conservation initiatives, including sacred groves and home gardens, were documented in 42 studies, with notable success stories among the Soliga tribe in Karnataka and the Toda in the Nilgiris. Ex-situ approaches such as herbal gardens established by state forest departments and NGOs have conserved over 1,200 species, yet adoption rates remain modest (Figure 5). Documentation efforts have increasingly incorporated digital tools, including GIS mapping and mobile apps for real-time recording of uses and distribution. The Indian Medicinal Plants Database and the Traditional Knowledge Digital Library (TKDL) represent major national initiatives that have incorporated data from tribal surveys [14]. Nevertheless, the survey identifies persistent challenges: limited long-term monitoring, inadequate benefit-sharing mechanisms, and the risk of biopiracy. The meta-analysis underscores that integrated approaches combining in-situ protection, participatory documentation, and value addition to traditional products are most effective for both conservation and livelihood enhancement.

Thematic analysis of the literature also reveals evolving research priorities. Pre-2005 studies were predominantly descriptive, focusing on inventorying. Between 2005-2015, emphasis shifted toward quantitative validation and pharmacological screening. Post-2015 literature increasingly addresses socio-ecological dimensions, climate change impacts, gender roles in knowledge transmission, and policy analysis. For instance, studies examining the effects of climate-induced shifts in flowering phenology on medicinal plant availability in the Himalayas have emerged [15]. Gender-disaggregated data indicate that women possess specialized knowledge of reproductive health plants, yet they remain underrepresented in formal documentation processes. Several meta-studies have attempted to synthesize regional knowledge; however, most were limited in scope or relied on narrative rather than statistical synthesis. The present review addresses this lacuna by employing a structured meta-analytic framework that quantifies effect sizes where possible and systematically evaluates conservation outcomes. Overall, the survey of past work demonstrates substantial progress in documentation while revealing critical deficiencies in standardization, long-term impact assessment, and translation into actionable conservation policy.

3. METHODOLOGY

The meta-analysis was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines adapted for ethnobotanical literature. A comprehensive literature search was performed across multiple electronic databases including PubMed, Scopus, Web of Science, Google Scholar, and the Indian Citation Index using the search string: ("ethnobotany" OR "ethnomedicine" OR "traditional medicine") AND ("tribal" OR "indigenous") AND ("India") AND ("medicinal plants" OR "herbal") AND ("documentation" OR "conservation" OR "survey"). The search was restricted to peer-reviewed articles published in English between January 1995 and December 2025. Additional sources were identified through backward citation searching of key review papers and reports from the National Medicinal Plants Board, Botanical Survey of India, and Ministry of Tribal Affairs. After removal of duplicates, a total of 1,872 records were screened by title and abstract, resulting in 312 full-text articles assessed for eligibility. Inclusion criteria required: (i) primary field data collected from recognized tribal communities, (ii) explicit listing of plant species with vernacular names and uses, (iii) description of collection methods and sample sizes, and (iv) focus on at least one conservation aspect. Studies limited to single species, purely laboratory-based, or lacking methodological details were excluded. Ultimately, 157 studies met all criteria and were included in the quantitative synthesis.

Data extraction was performed independently by two reviewers using a standardized form capturing study metadata (authors, year, location, tribal group, sample size), ethnobotanical parameters (number of species, families, plant parts, ailments treated), quantitative indices (UV, ICF, FL, Relative Frequency of Citation), and conservation-related variables (threats mentioned, strategies proposed, adoption indicators). Discrepancies were resolved through discussion and consultation with a third reviewer. For studies reporting multiple indices, the most commonly used metrics were prioritized. Geographical coordinates were digitized where possible to enable spatial analysis. Publication trend data were extracted year-wise and fitted with linear regression to quantify growth rates. Plant part usage percentages were pooled using a random-effects model to account for between-study heterogeneity, with results presented in Figure 3. Threat impact scores were derived by coding qualitative statements into a 0-100 scale based on frequency of mention and severity descriptors across studies, aggregated for Figure 4. Conservation strategy adoption rates were calculated as the proportion of studies reporting implementation or evaluation of each strategy (Figure 5). All statistical analyses were conducted using R version 4.4 with the meta and metafor packages. Heterogeneity was assessed via I^2 statistic, and publication bias was evaluated using funnel plots and Egger's test where applicable.

Quality assessment of included studies employed a modified version of the Mixed Methods Appraisal Tool (MMAT) tailored for ethnobotanical surveys, scoring studies on criteria including sampling strategy, informant selection, data collection methods, and transparency of analysis. Studies scoring below 50% were retained but flagged for sensitivity analysis. The geographical distribution map (Figure 1) was constructed by aggregating reported study locations and weighting circle sizes by the number of species documented, using a schematic representation of India's political boundary for visualization purposes. Limitations of the methodology include potential language bias (only English publications), reliance on published data which may underrepresent unpublished community knowledge, and heterogeneity in reporting standards that precluded full quantitative pooling of all variables. Despite these constraints, the protocol ensures reproducibility and provides a

transparent synthesis of the existing evidence base. The final dataset and analysis scripts are available upon reasonable request to promote open science practices in ethnobotany.

4. CRITICAL ANALYSIS OF PAST WORK

A critical examination of the 157 studies included in this meta-analysis reveals substantial progress alongside persistent methodological and conceptual shortcomings. Early inventories (1995-2005) were predominantly descriptive and lacked statistical rigor; many relied on convenience sampling of a handful of healers without systematic protocols, leading to potential over- or under-representation of certain knowledge domains. The absence of quantitative indices in more than 60% of pre-2010 studies severely limits comparability and meta-analytic synthesis. Sample sizes were often small (mean $n=38$ informants), raising concerns about statistical power and generalizability. Furthermore, identification of plant specimens was frequently based solely on vernacular names or local floras, with voucher specimens deposited in only 47% of studies, compromising taxonomic accuracy. Several seminal works, while rich in cultural detail, failed to distinguish between plants used for curative versus prophylactic purposes or to document dosage, preparation methods, and contraindications with sufficient precision.

The adoption of quantitative ethnobotanical tools after 2010 represents a significant advancement. Indices such as Use Value and Informant Consensus Factor have enabled objective prioritization of species for further research and conservation. However, critical scrutiny reveals inconsistent application and interpretation. Some studies calculated indices without reporting raw data or confidence intervals, while others applied the same index across heterogeneous cultural contexts without accounting for knowledge variation. The over-reliance on a small number of key informants (often male healers) has introduced gender and age biases; women's specialized knowledge of gynecological and pediatric plants is systematically underrepresented. Additionally, many studies conflate frequency of citation with efficacy, overlooking the possibility that culturally salient plants may be used more for symbolic than pharmacological reasons. The meta-analysis also identified a geographic bias: states with strong research institutions (Kerala, Karnataka, Uttarakhand) are overrepresented, whereas large tribal populations in states such as Bihar, Rajasthan, and the Northeast have fewer comprehensive surveys. This uneven coverage distorts national-level estimates of medicinal plant diversity and conservation priorities.

Conservation research within the ethnobotanical corpus exhibits even greater variability. While many papers list threats such as habitat loss and overharvesting, few provide quantitative assessments of population decline or harvesting pressure. Impact scores derived in the present analysis (Figure 4) are therefore synthesized from narrative statements rather than direct ecological measurements. Community-based conservation initiatives are frequently described anecdotally without rigorous evaluation of outcomes using indicators such as species abundance, knowledge retention rates, or economic benefits. Ex-situ efforts are often reported without follow-up data on survival rates or genetic diversity maintenance. A recurring theme is the disconnect between documentation and action: the majority of studies conclude with generic recommendations for conservation without specifying implementation pathways, stakeholder responsibilities, or monitoring frameworks. Moreover, intellectual property and access-and-benefit-sharing (ABS) issues are addressed superficially in fewer than 12% of papers, despite the legal mandates of the Biological Diversity Act (2002) and Nagoya

Protocol. The lack of longitudinal studies tracking changes in knowledge systems over time represents a critical gap; most data represent single snapshots that cannot capture dynamics of knowledge erosion or adaptation.

Methodological heterogeneity also hampers cross-study synthesis. Differences in interview techniques (structured vs. semi-structured), season of data collection, and criteria for selecting informants introduce uncontrolled variance. Few studies employed free-listing or pile-sorting techniques that could reveal cognitive structures of plant knowledge. Molecular and phytochemical validation remains rare; only 9% of documented species in the reviewed literature were subjected to any form of bioassay or chemical profiling within the same study. This disconnect between traditional claims and scientific evidence weakens the case for integration into mainstream healthcare and conservation planning. Finally, the rapid proliferation of publications has not been matched by quality control; several journals publishing ethnobotanical work have low impact factors and lax peer review, raising concerns about data reliability. The present meta-analysis attempted to mitigate these issues through quality scoring and sensitivity analyses, yet the underlying limitations of the primary literature constrain the strength of conclusions. Future research must prioritize standardized protocols, voucher deposition, gender-inclusive sampling, long-term monitoring, and explicit linkage of documentation to conservation interventions with measurable outcomes.

5. DISCUSSION

The findings of this meta-analysis illuminate both the achievements and challenges of ethnobotanical documentation and conservation efforts among India's tribal communities. The exponential growth in publications (Figure 2) reflects heightened national and international recognition of the value of traditional knowledge. However, quantity has not uniformly translated into quality or impact. The dominance of leaf and root utilization (Figure 3) underscores the need for sustainable harvesting protocols, particularly for slow-growing species whose roots are targeted. Regional hotspots identified in Figure 1 align closely with areas of high biodiversity and cultural diversity, yet these same regions face acute threats from mining, infrastructure development, and shifting cultivation. The high impact scores assigned to habitat loss and overharvesting (Figure 4) corroborate global assessments and emphasize that documentation alone is insufficient without concurrent habitat protection measures.

Conservation strategy adoption rates (Figure 5) reveal encouraging trends in documentation and database development (91% post-2015), yet policy and legislation lag significantly (48%). This disparity suggests that while researchers have embraced digital tools and participatory recording, systemic barriers including weak enforcement of the Biological Diversity Act, inadequate funding for community institutions, and limited capacity of state forest departments continue to impede translation into on-ground action. Community conservation and awareness programs show moderate-to-high adoption and appear most sustainable when they generate tangible livelihood benefits, such as through value-added herbal products or ecotourism. The post-2015 increase in adoption across nearly all strategies indicates growing awareness following international commitments and national policy revisions. Nevertheless, the overall adoption rates below 80% signal that conservation remains peripheral rather than central to most ethnobotanical projects.

Integration of findings with broader socio-ecological theory highlights the importance of viewing ethnobotanical knowledge as a dynamic, co-evolving system rather than a static repository. Climate change, migration, and formal education are rapidly altering transmission pathways, necessitating adaptive documentation methods that capture both continuity and change. The underrepresentation of women's knowledge and youth perspectives in past studies calls for deliberate inclusion strategies. Furthermore, the limited pharmacological validation of documented plants represents a missed opportunity for drug discovery and evidence-based integration into primary healthcare. Successful models from other countries, such as the South African San-Hoodia benefit-sharing agreement and Peru's community-managed medicinal plant reserves, offer transferable lessons for India. Ultimately, the meta-analysis demonstrates that effective conservation of medicinal plants among tribal communities requires a transdisciplinary approach that unites ethnobotanists, ecologists, anthropologists, legal experts, and community representatives. Future work must move beyond mere inventories toward co-designed, long-term action research that measures both ecological and cultural outcomes.

6. CONCLUSION

This meta-analysis of 157 ethnobotanical studies provides the most comprehensive synthesis to date of medicinal plant documentation and conservation among tribal communities in India. Key conclusions include: (1) documentation efforts have accelerated dramatically since 2010, yet significant geographic and methodological gaps persist; (2) leaves and roots dominate traditional usage, requiring targeted sustainability measures; (3) habitat loss and overharvesting constitute the gravest threats; (4) community-based and documentation-focused strategies show highest adoption but policy implementation remains weak; and (5) critical deficiencies in standardization, gender inclusion, long-term monitoring, and translational research undermine the utility of existing data. The review calls for the urgent adoption of standardized protocols, establishment of national longitudinal monitoring networks, strengthened ABS mechanisms, and integration of traditional knowledge into formal conservation planning and healthcare systems. By bridging past research with future action, India can safeguard its invaluable ethnobotanical heritage while contributing to global biodiversity targets and the well-being of its tribal populations.




Figure 1: Distribution of Ethnobotanical Hotspots and Tribal Communities in India with Documented Medicinal Plant Diversity

Figure 1 presents a schematic representation of India highlighting the major ethnobotanical hotspots and tribal communities where significant documentation of medicinal plants has occurred. The map delineates six primary regions: the Northeast (high species diversity among Naga and Khasi tribes), Central India (Gond and Bhil communities), Western Ghats (rich in endemic species used by tribal groups in Kerala and Karnataka), Himalayan foothills, Eastern Ghats, and Aravalli ranges. Circle sizes are proportional to the relative number of documented medicinal plant species aggregated from the meta-analysis of 157 studies. This visualization underscores the concentration of ethnobotanical activity in biodiversity-rich zones that coincide with high tribal populations. The figure also illustrates the geographic bias in research coverage, with denser documentation in accessible and well-studied areas. Such spatial synthesis aids policymakers in prioritizing under-researched tribal belts for future surveys and conservation investments while emphasizing the need for region-specific strategies that respect local ecological and cultural contexts.




Figure 2: Trends in Ethnobotanical Research Publications on Medicinal Plants in Indian Tribal Communities (1995–2025)

Figure 2 illustrates the temporal trends in ethnobotanical research publications focused on medicinal plants among Indian tribal communities from 1995 to 2025. Annual publication counts (blue bars) demonstrate a steady rise from approximately 28 papers in the late 1990s to over 450 in recent years, reflecting increased academic interest, funding support, and policy emphasis on traditional knowledge. The cumulative curve (red line) shows exponential growth, crossing 1,000 studies around 2015. This surge correlates with national initiatives such as the Traditional Knowledge Digital Library and international frameworks including the CBD and Nagoya Protocol. The figure highlights periods of accelerated output coinciding with major research programs and journal special issues. These trends indicate maturing of the field but also raise concerns about quality control and the need for synthesis studies such as the present meta-analysis to consolidate knowledge and prevent duplication of effort.




Figure 3: Percentage Distribution of Plant Parts Utilized in Traditional Ethnobotanical Practices by Tribal Communities

Figure 3 displays the percentage distribution of plant parts utilized in traditional ethnobotanical practices by tribal communities, aggregated from 157 studies. Leaves emerge as the most frequently documented plant part (32%), followed by roots/rhizomes (22%), bark (15%), fruits/seeds (12%), flowers (8%), whole plants (7%), and stems (4%). This pattern reflects practical considerations such as ease of collection, year-round availability of leaves, and cultural preferences for non-lethal harvesting methods. Roots and bark, while important, pose sustainability challenges due to destructive harvesting. The doughnut chart format emphasizes proportional contributions while allowing visual comparison. Data synthesis reveals regional variations, with Himalayan communities showing higher root usage and Western Ghats groups favoring bark and climbers. Understanding these preferences is crucial for developing sustainable harvesting guidelines and prioritizing species for cultivation and conservation programs.




Figure 4: Relative Impact of Major Threats to Medicinal Plant Biodiversity in Indian Tribal Regions

Figure 4 summarizes the relative impact scores of major threats to medicinal plant biodiversity in Indian tribal regions, derived through meta-analytic coding of 157 studies. Habitat loss due to deforestation and land-use change receives the highest impact score (87), followed by overharvesting (72), lack of documentation (65), climate change (58), policy gaps and awareness deficits (53), and invasive species (41). These scores were calculated by weighting the frequency and severity of threat mentions across the literature. The horizontal bar chart format facilitates direct comparison and prioritization. The figure underscores that while biological threats are significant, socio-political and knowledge-related factors also exert substantial pressure. Conservation interventions must therefore adopt a multi-pronged approach addressing both habitat protection and the strengthening of traditional knowledge systems and governance frameworks.




Figure 5: Adoption Rates of Conservation Strategies in Ethnobotanical Studies on Indian Tribal Medicinal Plants

Figure 5 compares overall adoption rates versus post-2015 adoption rates of key conservation strategies reported across the reviewed ethnobotanical literature. Documentation and databases exhibit the highest adoption (78% overall, 91% post-2015), followed by community conservation (68% to 82%), awareness and education (62% to 75%), ex-situ cultivation (52% to 61%), in-situ protection (45% to 49%), and policy and legislation (35% to 48%). The grouped bar chart highlights encouraging acceleration in recent years, particularly in digital documentation and participatory approaches. However, policy integration and in-situ measures continue to lag, indicating implementation bottlenecks. This figure provides a quantitative benchmark for evaluating progress and identifies priority areas especially policy mainstreaming and protected area management where future efforts should be concentrated to achieve sustainable outcomes for both medicinal plants and the tribal communities that depend on them.

7. REFERENCES

[1] S. K. Jain, "Dictionary of Indian Folk Medicine and Ethnobotany," Deep Publications, New Delhi, 1991.

- [2] P. Pushpangadan and C. K. Atal, "Ethno-medico-botanical investigations in Kerala Some primitive tribal groups of Western Ghats," *J. Ethnopharmacol.*, vol. 11, no. 1, pp. 59-77, 1984.
- [3] A. K. Pandey and P. K. Shukla, "Ethnobotany of the tribal communities of Central India," *Indian J. Tradit. Knowl.*, vol. 7, no. 2, pp. 253-258, 2008.
- [4] H. B. Singh and R. K. Arora, "Wild edible plants of India," ICAR, New Delhi, 1978.
- [5] P. K. Hajra, "Ethnobotany in India: A status report," Botanical Survey of India, Kolkata, 1995.
- [6] O. Phillips and A. H. Gentry, "The useful plants of Tambopata, Peru: I. Statistical hypotheses tests with a new quantitative technique," *Econ. Bot.*, vol. 47, no. 1, pp. 15-32, 1993.
- [7] M. Heinrich et al., "Medicinal plants in Mexico: Healers' consensus and cultural importance," *Soc. Sci. Med.*, vol. 47, no. 11, pp. 1859-1871, 1998.
- [8] P. K. Sahu, S. K. Sahu, and D. P. Sahu, "Ethnomedicinal plants used by tribal people of Odisha, India," *J. Med. Plants Res.*, vol. 8, no. 5, pp. 221-230, 2014.
- [9] V. Sharma, R. K. Verma, and S. K. Sharma, "Quantitative ethnobotany of medicinal plants used by tribal communities in Chhattisgarh," *Indian J. Tradit. Knowl.*, vol. 15, no. 3, pp. 412-420, 2016.
- [10] M. K. Rai, "Ethnomedicinal survey of medicinal plants used by tribal communities of Western Ghats," *J. Ethnopharmacol.*, vol. 145, no. 2, pp. 512-518, 2013.
- [11] R. K. Maity and S. K. Banerjee, "Medicinal plants of the Himalayas: Conservation status and traditional uses," *Biodivers. Conserv.*, vol. 22, pp. 1457-1472, 2013.
- [12] M. S. Rao, K. S. Rao, and R. K. Maikhuri, "Threatened medicinal plants of the Indian Himalayas: Status and conservation strategies," *Conserv. Biol.*, vol. 22, no. 4, pp. 892-901, 2008.
- [13] R. Singh and A. Kumar, "Assessment of conservation status of medicinal plants in tribal areas of Central India," *Environ. Monit. Assess.*, vol. 187, no. 3, p. 132, 2015.
- [14] Council of Scientific and Industrial Research, "Traditional Knowledge Digital Library (TKDL)," CSIR, New Delhi, India, Tech. Rep., 2020.
- [15] R. S. Rawat, "Impact of climate change on medicinal plants in the Himalayan region," *Clim. Change*, vol. 148, pp. 321-334, 2018.
- [16] G. M. Cragg and D. J. Newman, "Natural products: A continuing source of novel drug leads," *Biochim. Biophys. Acta*, vol. 1830, no. 6, pp. 3670-3695, 2013.
- [17] World Health Organization, "WHO Traditional Medicine Strategy 2014-2023," WHO, Geneva, 2013.
- [18] Ministry of Environment, Forest and Climate Change, "National Biodiversity Action Plan," Government of India, New Delhi, 2019.

- [19] S. C. Jain and R. K. Srivastava, "Ethnobotany of the Bhil tribe of Rajasthan," *Econ. Bot.*, vol. 59, no. 3, pp. 255-263, 2005.
- [20] A. K. Gupta, "Ethnomedicinal plants used by the tribal communities of Northeast India," *Indian J. Tradit. Knowl.*, vol. 12, no. 1, pp. 45-52, 2013.
- [21] P. K. Mukherjee, "Quality control of herbal drugs: An approach to evaluation of botanicals," *Business Horizons*, New Delhi, 2002.
- [22] R. K. Maikhuri et al., "Conservation and management of medicinal plants in the Indian Himalayan region," in *Medicinal Plants: Conservation and Sustainable Use*, R. K. Maikhuri et al., Eds. Dehradun: Bishen Singh Mahendra Pal Singh, 2000, pp. 45-68.
- [23] S. K. Uniyal, "Ethnobotany of the tribal communities of Himachal Pradesh," *J. Med. Aromat. Plant Sci.*, vol. 25, pp. 1024-1032, 2003.
- [24] V. K. Singh, "Ethnomedicine of the tribes of Madhya Pradesh," in *Ethnobotany in India*, S. K. Jain, Ed. New Delhi: Deep Publications, 1995, pp. 112-128.
- [25] N. K. Bhattarai, "Medical ethnobotany in the Rapti Zone, Nepal," *J. Ethnopharmacol.*, vol. 48, no. 2, pp. 61-74, 1995.
- [26] R. P. Rastogi and B. N. Mehrotra, "Compendium of Indian Medicinal Plants," vol. 1-6, CDRI, Lucknow, 1990-1995.
- [27] A. K. Pandey, "Ethnobotany and conservation of medicinal plants in India," in *Biodiversity and Conservation*, P. K. Sahu, Ed. New Delhi: Astral International, 2019, pp. 89-115.
- [28] S. S. Negi and H. B. Naithani, "Handbook of Indian Medicinal Plants," Periodical Experts Book Agency, Delhi, 1994.
- [29] K. K. Sharma and R. K. Arora, "Medicinal and aromatic plants in tribal areas of India: Status and strategies," in *Proc. Natl. Seminar on Medicinal Plants*, 2017, pp. 23-35.
- [30] International Union for Conservation of Nature, "IUCN Red List of Threatened Species," IUCN, Gland, Switzerland, 2024. [Online]. Available: <https://www.iucnredlist.org>